




[Home](#) [Publications](#) [Resources](#) [Librarians](#) [Press](#) [Advertise](#)

 Hello. Sign in to personalize your visit. New user? [Register now](#).

Author information

Nabihah Haron, BSc,¹ Rosnah Binti Zain, MSc,^{2,3} Wan Maria Nabillah, MMedSc,² Amyza Saleh, PhD,¹ Thomas George Kallarakkal, MDS,^{2,3} Anand Ramanathan, MDS,^{2,3} Suraya Hani Mohd Sinon, DClinDent,⁴ Ishak Abdul Razak, PhD,^{3,5} and Sok Ching Cheong, PhD^{2,6}

¹Outreach Team, Cancer Research Malaysia, Subang Jaya, Malaysia.

²Department of Oral and Maxillofacial Clinical Sciences, Faculty of Dentistry, University of Malaya, Kuala Lumpur, Malaysia.

³Oral Cancer Research and Coordinating Centre (OCRCC), University of Malaya, Kuala Lumpur, Malaysia.

⁴Department of Oral Pathology and Oral Medicine, Faculty of Dentistry, University Kebangsaan Malaysia, Kuala Lumpur, Malaysia.

⁵Department of Community Oral Health & Clinical Prevention, Faculty of Dentistry, University of Malaya, Kuala Lumpur, Malaysia.

⁶Oral Cancer Research Team, Cancer Research Malaysia, Subang Jaya, Malaysia.

Address correspondence to:

Sok Ching Cheong, PhD

Oral Cancer Research Team

Cancer Research Malaysia

No. 1, Jalan SS12/1A

Subang Jaya

Selangor 47500

Malaysia

E-mail: sokching.cheong@cancerresearch.my

Received: June 8, 2016

Revised: July 10, 2016

Accepted: July 11, 2016

ABSTRACT

Introduction: This study examined the concordance in clinical diagnosis of high-risk lesions in the oral cavity and referral decisions between clinical oral examination (COE) and teledentistry. **Materials and Methods:** Sixteen individuals with a range of oral potentially malignant disorders (OPMD) and normal oral mucosa were included. Five areas of the oral cavity were photographed by three dentists using mobile phone cameras with 5 MP-13 MP resolutions. On the same day, the patients were given COE by two oral medicine specialists (OMS) and 3 weeks later, they reviewed the images taken using the phone, and concordance was examined between the two by Kappa statistics. The sensitivity and specificity of clinical diagnosis using the phone images were also measured. Pre- and post-program questionnaires were answered by both the dentists and the OMS to determine the feasibility of integrating teledentistry in their clinical practice. **Results:** The Kappa values in determining the presence of lesion, category of lesion (OPMD or not), and making referral decision were moderate to strong (0.64–1.00). The overall sensitivity was more than 70% and specificity was 100%. The false negative rate decreased as the camera resolution increased. All dentists agreed that the process could facilitate early detection of oral mucosal lesion, and was easy to use in the clinic. **Conclusions:** This study provides evidence that teledentistry can be used for communication between primary care and OMS and could be readily integrated into clinical setting for patient management.

You have requested the following:



Mobile Phone Imaging in Low Resource Settings for Early Detection of Oral Cancer and Concordance with Clinical Oral Examination

Haron Nabihah, Zain Rosnah Binti, Nabillah Wan Maria, Saleh Amyza, Kallarakkal Thomas George, Ramanathan Anand, Sinon Suraya Hani Mohd, Razak Ishak Abdul, and Cheong Sok Ching

Telemicine and e-Health. March 2017, Vol. 23, No. 3: 192-199

- PAY PER VIEW** Telemicine and e-Health - 23(3):192-199; Mobile Phone Imaging in Low Resource Settings for Early Detection of Oral Cancer and Concordance with Clinical Oral Examination (access for 24 hours for US \$51.00)

[ADD TO CART](#)

SUBSCRIBER LOGIN

Already a subscriber? [Login](#) to access this journal.